

Lamya H. Al-Wahaibi, Nora H. Al-Shaalan, Hazem A. Ghabbour, Edward R.T. Tiekink and Ali A. El-Emam*

Crystal structure of 3,5-bis(trifluoromethyl)benzyl (Z)-N'-(adamantan-1-yl)-4-phenylpiperazine-1-carbothioimide, C₃₀H₃₃F₆N₃S




Table 1: Data collection and handling.

Crystal:	Slab, colorless
Size:	0.58 × 0.47 × 0.07 mm
Wavelength:	Mo K α radiation (0.71073 Å)
μ :	0.18 mm ⁻¹
Diffractometer, scan mode:	Bruker APEX-II, φ and ω -scans
θ_{\max} , completeness:	33.2°, >99%
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	48368, 10750, 0.095
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 7424
$N(\text{param})_{\text{refined}}$:	398
Programs:	Bruker programs [1], SHELX [2, 3], ORTEP [4]

<https://doi.org/10.1515/ncrs-2019-0220>

Received March 24, 2019; accepted April 23, 2019; available online May 27, 2019

Abstract

C₃₀H₃₃F₆N₃S, triclinic, $P\bar{1}$ (no. 2), $a = 9.3012(5)$ Å, $b = 10.2734(5)$ Å, $c = 15.1850(8)$ Å, $\alpha = 81.982(2)^\circ$, $\beta = 78.696(2)^\circ$, $\gamma = 83.882(2)^\circ$, $V = 1404.25(13)$ Å³, $Z = 2$, $R_{\text{gt}}(F) = 0.0779$, $wR_{\text{ref}}(F^2) = 0.2499$, $T = 293(2)$ K.

CCDC no.: 1554490

The crystal structure is shown in the figure. Table 1 contains crystallographic data and Table 2 contains the list of the atoms including atomic coordinates and displacement parameters.

*Corresponding author: Ali A. El-Emam, Department of Medicinal Chemistry, Faculty of Pharmacy, University of Mansoura, Mansoura 35516, Egypt, e-mail: elemam5@hotmail.com

Lamya H. Al-Wahaibi and Nora H. Al-Shaalan: Department of Chemistry, College of Sciences, Princess Nourah Bint Abdulrahman University, Riyadh 11671, Saudi Arabia

Hazem A. Ghabbour: Department of Medicinal Chemistry, Faculty of Pharmacy, University of Mansoura, Mansoura 35516, Egypt

Edward R.T. Tiekink: Research Centre for Crystalline Materials, School of Science and Technology, Sunway University, 47500 Bandar Sunway, Selangor Darul Ehsan, Malaysia

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.98386(5)	0.51000(4)	0.66054(3)	0.01323(12)
N1	0.69696(17)	0.60257(15)	0.64402(11)	0.0122(3)
N2	0.86822(17)	0.75759(15)	0.62935(11)	0.0129(3)
N3	0.82652(17)	0.92552(15)	0.46515(11)	0.0141(3)
F1 ^a	0.6835(5)	0.8971(3)	1.02509(18)	0.0589(9)
F2 ^a	0.8013(4)	0.9811(2)	0.8977(2)	0.0610(9)
F3 ^a	0.5984(4)	0.9168(3)	0.9015(2)	0.0706(9)
F1 ^b	0.8490(9)	0.9393(6)	0.9780(5)	0.0619(13)
F2 ^b	0.7236(12)	0.9677(7)	0.8762(5)	0.0598(14)
F3 ^b	0.6193(10)	0.8858(9)	1.0038(6)	0.0622(15)
F4	0.72337(19)	0.41464(15)	1.10754(9)	0.0349(4)
F5	0.8472(2)	0.29227(15)	1.01309(12)	0.0573(6)
F6	0.6274(3)	0.3624(2)	1.00252(14)	0.0643(7)
C1	0.82839(19)	0.62775(17)	0.64263(12)	0.0116(3)
C2	0.9920(2)	0.79314(18)	0.55694(13)	0.0150(3)
H2A	1.0694	0.7219	0.5558	0.018*
H2B	1.0313	0.8711	0.5692	0.018*
C3	0.9452(2)	0.81997(18)	0.46506(13)	0.0144(3)
H3A	1.0287	0.8450	0.4188	0.017*
H3B	0.9121	0.7403	0.4508	0.017*
C4	0.7000(2)	0.89565(18)	0.53841(13)	0.0148(3)
H4A	0.6542	0.8212	0.5262	0.018*
H4B	0.6276	0.9710	0.5404	0.018*
C5	0.7496(2)	0.86352(18)	0.62956(13)	0.0146(3)
H5A	0.7829	0.9422	0.6452	0.017*

Table 2 (continued)

Atom	x	y	z	U_{iso}^*/U_{eq}
H5B	0.6663	0.8377	0.6757	0.017*
C6	0.7924(2)	0.97605(18)	0.37967(14)	0.0156(4)
C7	0.6890(2)	1.08575(19)	0.37243(16)	0.0200(4)
H7	0.6424	1.1225	0.4243	0.024*
C8	0.6557(2)	1.1396(2)	0.28977(17)	0.0253(5)
H8	0.5862	1.2114	0.2867	0.030*
C9	0.7252(3)	1.0876(2)	0.21042(17)	0.0281(5)
H9	0.7018	1.1236	0.1548	0.034*
C10	0.8297(3)	0.9814(2)	0.21635(16)	0.0262(5)
H10	0.8783	0.9469	0.1640	0.031*
C11	0.8626(2)	0.9256(2)	0.29992(15)	0.0201(4)
H11	0.9323	0.8539	0.3026	0.024*
C12	0.63842(19)	0.47279(17)	0.66558(12)	0.0105(3)
C13	0.47856(19)	0.49282(17)	0.65042(13)	0.0131(3)
H13A	0.4230	0.5544	0.6899	0.016*
H13B	0.4769	0.5306	0.5884	0.016*
C14	0.4059(2)	0.36227(18)	0.66950(13)	0.0142(3)
H14	0.3044	0.3778	0.6592	0.017*
C15	0.4064(2)	0.30470(19)	0.76833(13)	0.0162(4)
H15A	0.3500	0.3650	0.8084	0.019*
H15B	0.3611	0.2217	0.7812	0.019*
C16	0.5654(2)	0.28244(19)	0.78459(13)	0.0152(3)
H16	0.5656	0.2454	0.8476	0.018*
C17	0.6379(2)	0.41381(18)	0.76491(12)	0.0139(3)
H17A	0.7380	0.3991	0.7758	0.017*
H17B	0.5840	0.4752	0.8051	0.017*
C18	0.7227(2)	0.37406(17)	0.60209(12)	0.0127(3)
H18A	0.7217	0.4103	0.5397	0.015*
H18B	0.8243	0.3595	0.6102	0.015*
C19	0.6515(2)	0.24254(17)	0.62253(13)	0.0135(3)
H19	0.7073	0.1803	0.5825	0.016*
C20	0.4927(2)	0.26407(18)	0.60698(13)	0.0149(3)
H20A	0.4478	0.1809	0.6198	0.018*
H20B	0.4911	0.2984	0.5443	0.018*
C21	0.6524(2)	0.18559(18)	0.72137(13)	0.0159(4)
H21A	0.7529	0.1701	0.7315	0.019*
H21B	0.6084	0.1019	0.7342	0.019*
C22	1.0637(2)	0.5927(2)	0.73707(13)	0.0160(4)
H22A	1.1509	0.5404	0.7517	0.019*
H22B	1.0929	0.6780	0.7070	0.019*
C23	0.9554(2)	0.61106(19)	0.82258(13)	0.0150(3)
C24	0.8961(2)	0.7359(2)	0.84203(14)	0.0205(4)
H24	0.9264	0.8099	0.8029	0.025*
C25	0.7908(3)	0.7504(2)	0.92038(15)	0.0243(4)
C26	0.7426(3)	0.6411(2)	0.97887(14)	0.0230(4)
H26	0.6706	0.6509	1.0300	0.028*
C27	0.8041(2)	0.5168(2)	0.95945(14)	0.0192(4)
C28	0.9100(2)	0.50135(19)	0.88264(13)	0.0167(4)
H28	0.9510	0.4174	0.8711	0.020*
C29 ^a	0.7186(4)	0.8848(4)	0.9376(2)	0.0486(8)
C29 ^b	0.7479(9)	0.8848(11)	0.9475(5)	0.0535(10)
C30	0.7503(3)	0.3973(2)	1.02066(15)	0.0255(5)

Occupancies: ^a = 0.716(3), ^b = 284(3)

Source of material

3,5-Bis(trifluoromethyl)benzyl bromide (614 mg, 2.0 mmol) and anhydrous potassium carbonate (276 mg, 2 mmol) were added to an anhydrous acetone (15 mL) solution of *N*-(adamantan-1-yl)-4-phenylpiperazine-1-carbothioamide (711 mg, 2 mmol), and the mixture was heated under reflux for 4 h. The solvent was then distilled off *in vacuo* and the resulting residue was washed with water (20 mL), dried and crystallized from aqueous ethanol to yield 872 mg (75%) of the title compound as colourless crystals. *M.pt.*: 386–388 K. Single crystals suitable for X-ray analysis were obtained by the slow evaporation of its CHCl₃:EtOH (1:1; 5 mL) solution held at room temperature. ¹H NMR (CDCl₃, 700.17 MHz): δ 1.58 (s, 6H, adamantane-CH₂), 1.69 (s, 6H, adamantane-CH₂), 1.95–1.97 (m, 3H, adamantane-CH), 3.29–3.31 (m, 4H, piperazine-H), 3.40–3.42 (m, 4H, piperazine-H), 4.04 (s, 2H, benzylic CH₂), 6.93–7.02 (m, 3H, Ar-H), 7.29 (s, 1H, Ar-H), 7.32–7.34 (m, 2H, Ar-H), 7.78 (s, 2H, Ar-H). ¹³C{¹H} NMR (CDCl₃, 176.08 MHz): δ 29.80, 35.51, 36.44, 54.80 (adamantane-C), 37.31 (benzylic CH₂), 49.0, 49.13 (piperazine-C), 116.28, 120.12, 129.16, 129.24, 131.44, 131.63, 141.30, 147.50 (Ar-C), 124.04 (CF₃), 151.19 (C=N). **ESI-MS**, *m/z*: 582.2 [*M* + *H*]⁺.

Experimental details

The C-bound H atoms were geometrically placed (C–H = 0.93–0.98 Å) and refined as riding with $U_{iso}(H) = 1.2U_{eq}(C)$. Disorder was resolved for the C25-bound CF₃ group with two distinct sites modelled. The site occupancy factor for the major component refined to 0.716(3). Standard restraints in SHELXL [3], such as SADI and SIMU, were employed. The maximum and minimum residual electron density peaks of 1.14 and 1.35 e Å⁻³, respectively, were located 0.80 and 0.73 Å from the F1' and S1 atoms, respectively.

Discussion

Recently, a series of adamantane-isothiourea hybrid derivatives were synthesized and demonstrated to exhibit a broad spectrum of anti-bacterial activity [5]. The motivation for the synthesis of these potential drugs was twofold. Firstly, the adamantane cage, known to be highly lipophilic, is a crucial pharmacophore in a number of drugs [6]. A key example is that of the drug amantadine, an effective antiviral agent for influenza A viruses [7] as well as being a potent anti-parkinsonian agent [8]. Subsequently, adamantane derivatives have been shown to exhibit diversified pharmacological potential, including anti-viral [9], anti-cancer [10], anti-bacterial and anti-fungal [11], anti-malarial [12] and anti-diabetic [13] activities. Secondly, and in the same way, isothiourea derivatives are known to display

significant anti-viral [14], anti-cancer [15] and anti-bacterial [16] activities. With such a diverse range of pharmacological profiles exhibited by both adamantane and isothiourea derivatives, it was thought their combination through the synthesis of adamantane-isothiourea hybrid molecules would lead to even more efficient therapeutic agents [5]. Herein, the crystal and molecular structures of the title compound are described in continuation of complementary structural studies of these new anti-bacterial agents [17].

The molecule [systematic name: 3,5-bis(trifluoromethyl)benzyl 4-phenyl-*N*-(tricyclo[3.3.1.1^{3,7}]decan-1-yl)piperazine-1-carboximidothioate] is shown in the figure (70% displacement ellipsoids; the minor component of the disordered CF₃ has been omitted). The central CN₂S chromophore is strictly planar with the r.m.s. deviation of the least-squares plane through the C1, N1, N2 and S1 atoms being 0.0068 Å. There is a *Z* configuration about the C1=N1 bond [1.272(2) Å], and the piperazinyl ring has a chair conformation. The conformational relationship between the central plane and the piperazinyl ring approaches orthogonal with the dihedral angle between the least-squares planes through the residues being 70.42(7)°. The dihedral angle between the least-squares planes through the piperazinyl and the N3-bound phenyl ring is 28.47(6)°, indicating a twisted conformation. To a first approximation, the di-substituted benzyl residue is folded back over the rest of the molecule. Nevertheless, the residue is inclined towards the piperazinyl group as reflected in the values of the C22–S1–C1–N2 and C22–S1–C1–N1 torsion angles of 42.75(15) and –135.09(17)°, respectively. Generally the geometric parameters are in the expected ranges [18, 19].

The most prominent feature of the molecular packing are weak methylene-C–H···N(piperazinyl) interactions between centrosymmetrically related molecules [C2–H2b···N3ⁱ: H2b···N3ⁱ = 2.53 Å, C2···N3ⁱ = 3.444(2) Å with an angle at H2b = 157° for symmetry operation *i*: 2 – *x*, 2 – *y*, 1 – *z*], leading to dimeric aggregates. There are no other directional interactions connecting molecules. Globally, molecules assemble into double layers that stack along the *c* axis direction. The CF₃ groups project to either side of the layers and inter-digitate with successive layers. Reflecting the packing, the calculated Hirshfeld surfaces [20, 21] show a dominance of H···H [48.6%], F···H/H···F [28.1%], C···H/H···C [10.2%] and F···F [6.3%] contacts to the overall surface.

There are two literature precedents for the title compound, namely derivatives with 4-nitrobenzyl [5] and 4-bromobenzyl [22] groups. There are non-trivial differences in the conformations in these molecules. While there is nearly an orthogonal relationship between the central plane and the piperazinyl ring in the title structure, these residues are closer to co-planar in the literature structures, with comparable dihedral angles of 32.2(3) and 34.78(7)°, respectively.

Acknowledgements: This work was funded by the Deanship of Scientific Research at Princess Nourah bint Abdulrahman University through the Research Group Program (Grant No. RGP-1438–0010).

References

1. Bruker. SADABS, APEX2 and SAINT. Bruker AXS Inc., Madison, WI, USA (2014).
2. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr.* **A64** (2008) 112–122.
3. Sheldrick, G. M.: Crystal structure refinement with SHELXL. *Acta Crystallogr.* **C71** (2015) 3–8.
4. Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Cryst.* **45** (2012) 849–854.
5. Al-Wahaibi, L. H.; Hassan, H. M.; Abo-Kamar, A. M.; Ghabbour, H. A.; El-Emam, A. A.: Adamantane-isothiourea hybrid derivatives: Synthesis, characterization, *in vitro* antimicrobial and *in vivo* hypoglycemic activities. *Molecules* **22** (2017) 710–722.
6. Wanka, L.; Iqbal, K.; Schreiner, P. R.: The lipophilic bullet hits the targets: Medicinal chemistry of adamantane derivatives. *Chem. Rev.* **113** (2013) 3516–3604.
7. Togo, Y.; Hornick, R. B.; Dawkins, A. T.: Studies on induced influenza in man. I. Double blind studies designed to assess prophylactic efficacy of amantadine hydrochloride against A2/Rockville/1/65 strain. *J. Am. Med. Assoc.* **203** (1968) 1089–1094.
8. Schwab, R. S.; England Jr, A. C.; Poskanzer, D. C.; Young, R. R.: Amantadine in the treatment of Parkinson's disease. *J. Am. Med. Assoc.* **208** (1969) 1168–1170.
9. El-Emam, A. A.; Al-Deeb, O. A.; Al-Omar, M. A.; Lehmann, J.: Synthesis, antimicrobial, and anti-HIV-1 activity of certain 5-(1-adamantyl)-2-substituted thio-1,3,4-oxadiazoles and 5-(1-adamantyl)-3-substituted aminomethyl-1,3,4-oxadiazoline-2-thiones. *Bioorg. Med. Chem.* **12** (2004) 5107–5113.
10. Lorenzo, P.; Alvarez, R.; Ortiz, M. A.; Alvarez, S.; Piedrafita, F. J.; de Lera, Á. R.: Inhibition of IκB kinase-β and anticancer activities of novel chalcone adamantyl arotinoids. *J. Med. Chem.* **81** (2008) 5431–5440.
11. El-Emam, A. A.; Al-Tamimi, A.-M. S.; Al-Omar, M. A.; Alrashood, K. A.; Habib, E. E.: Synthesis and antimicrobial activity of novel 5-(1-adamantyl)-2-aminomethyl-4-substituted-1,2,4-triazoline-3-thiones. *Eur. J. Med. Chem.* **68** (2013) 96–102.
12. Dong, Y.; Wittlin, S.; Sriraghavan, K.; Chollet, J.; Charman, S. A.; Charman, W. N.; Scheurer, C.; Urwyler, H.; Tomas, J. S.; Snyder, C.; Creek, D. J.; Morizzi, J.; Koltun, M.; Matile, H.; Wang, X.; Padmanilayam, M.; Tang, Y.; Dorn, A.; Brun, R.; Vennerstrom, J. L.: The structure-activity relationship of the antimalarial ozonide arterolane (OZ277). *J. Med. Chem.* **53** (2010) 481–491.
13. Augeri, D. J.; Robl, J. A.; Betebenner, D. A.; Magnin, D. R.; Khanna, A.; Robertson, J. G.; Wang, A.; Simpkins, L. M.; Taunk, P.; Huang, Q.; Han, S.; Abboa-Offei, B.; Cap, M.; Xin, L.; Tao, L.; Tozzo, E.; Welzel, G. E.; Egan, D. M.; Marcinkeviciene, J.; Chang, S. Y.; Biller, S. A.; Kirby, M. S.; Parker, R. A.; Hamann, L. G.: Discovery and preclinical profile of saxagliptin (BMS-477118): A highly potent, long-acting, orally

- active dipeptidyl peptidase IV inhibitor for the treatment of type 2 diabetes. *J. Med. Chem.* **48** (2005) 5025–5037.
14. Thoma, G.; Streiff, M. B.; Kovarik, J.; Glickman, F.; Wagner, T.; Beerli, C.; Zerwes, H.: Orally bioavailable isothioureas block function of the chemokine receptor CXCR4 *in vitro* and *in vivo*. *J. Med. Chem.* **51** (2008) 7915–7920.
 15. Koronkiewicz, M.; Romiszewska, A.; Chilmonec, Z.; Kazimierzczuk, Z.: New benzimidazole-derived isothioureas as potential antileukemic agents – Studies *in vitro*. *Med. Chem.* **11** (2015) 364–372.
 16. Nicholson, A.; Perry, J. D.; James, A. L.; Stanforth, S. P.; Carnell, S.; Wilkinson, K.; Anjam Khan, C. M.; De Soyza, A.; Gould, F. K.: *In vitro* activity of S-(3,4-dichlorobenzyl) isothiourea hydrochloride and novel structurally related compounds against multidrug-resistant bacteria, including *Pseudomonas aeruginosa* and *Burkholderia cepacia* complex. *Int. J. Antimicrob. Agents* **39** (2012) 27–32.
 17. Al-Ghulikah, H. A.; Ghabbour, H. A.; Tiekink, E. R. T.; El-Emam, A. A.: Crystal structure of 4-bromobenzyl (Z)-N-(adamantan-1-yl)morpholine-4-carbothioimide, C₂₂H₂₉BrN₂OS. *Z. Kristallogr. NCS* **234** (2019) “In Press” (<https://doi.org/10.1515/nrcs-2019-0216>).
 18. Al-Omary, F. A. M.; Alanazi, F. S.; Ghabbour, H. A.; El-Emam, A. A.: Crystal structure of 2-[3,5-bis(trifluoromethyl)benzylsulfanyl]-5-(5-bromothiophen-2-yl)-1,3,4-oxadiazole, C₁₅H₇BrF₆N₂OS₂. *Z. Kristallogr. NCS* **232** (2017) 131–133.
 19. Al-Wahaibi, L. H.; Hassan, H. M.; Ghabbour, H. A.; El-Emam, A. A.: Crystal structure of 3,5-bis(trifluoromethyl)benzyl(Z)-N-(adamantan-1-yl)morpholine-4-carbothioimide, C₂₄H₂₈F₆N₂OS. *Z. Kristallogr. NCS* **233** (2018) 607–609.
 20. Turner, M. J.; Mckinnon, J. J.; Wolff, S. K.; Grimwood, D. J.; Spackman, P. R.; Jayatilaka, D.; Spackman, M. A.: Crystal Explorer v17. The University of Western Australia, Australia (2017).
 21. Tan, S. L.; Jotani, M. M.; Tiekink, E. R. T.: Utilizing Hirshfeld surface calculations, non-covalent interaction (NCI) plots and the calculation of interaction energies in the analysis of molecular packing. *Acta Crystallogr.* **E75** (2019) 308–318.
 22. Al-Wahaibi, L. H.; Hassan, H. M.; Abo-Kamar, A. M.; Ghabbour, H. A.; El-Emam, A. A.: Crystal structure of 4-bromobenzyl (Z)-N'-(adamantan-1-yl)-4-phenylpiperazine-1-carbothioimide, C₂₈H₃₄BrN₃S. *Z. Kristallogr. NCS* **232** (2017) 189–189.