

See Mun Lee, Kong Mun Lo and Edward R.T. Tiekink*

Crystal structure of bis[(μ_3 -oxido)-(μ_2 -(*N,N*-diisopropylthiocarbamoylthio)acetato- $\kappa^2 O, O'$)-((*N,N*-diisopropylthiocarbamoylthio)acetato- κO)-bis(di-4-methylbenzyl-tin(IV))], $C_{100}H_{136}N_4O_{10}S_8Sn_4$




Table 1: Data collection and handling.

Crystal:	Colourless prism
Size:	0.30 × 0.20 × 0.06 mm
Wavelength:	Mo K α radiation (0.71073 Å)
μ :	1.13 mm ⁻¹
Diffractometer, scan mode:	CCD, φ and ω
θ_{\max} , completeness:	28.3°, >99%
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	25986, 13129, 0.033
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 10218
$N(\text{param})_{\text{refined}}$:	580
Programs:	Bruker [1], SHELX [2–4], WinGX/ORTEP [5]

<https://doi.org/10.1515/ncrs-2019-0161>

Received March 3, 2019; accepted March 25, 2019; available online June 19, 2019

Abstract

$C_{100}H_{136}N_4O_{10}S_8Sn_4$, triclinic, $P\bar{1}$ (no. 2), $a = 13.7756(4)$ Å, $b = 13.9663(5)$ Å, $c = 16.6392(5)$ Å, $\alpha = 71.501(2)^\circ$, $\beta = 73.952(2)^\circ$, $\gamma = 63.180(2)^\circ$, $V = 2675.49(16)$ Å³, $Z = 1$, $R_{\text{gt}}(F) = 0.0383$, $wR_{\text{ref}}(F^2) = 0.1209$, $T = 296(2)$ K.

CCDC no.: 1905385

The molecular structure is shown in the figure (hydrogen atoms are omitted for clarity). Table 1 contains crystallographic data and Table 2 contains the list of the atoms including atomic coordinates and displacement parameters.

Source of material

The melting point (uncorrected) of the compound was measured on an electrothermal digital melting point

apparatus. The elemental analysis was performed on a Perkin-Elmer EA2400 CHN analyser. The IR spectrum was recorded using a Perkin-Elmer RX1 spectrometer in a Nujol mull between KBr plates.

Di(4-methylbenzyl)tin dichloride was synthesized by the direct reaction of 4-methylbenzyl chloride (Sigma-Aldrich) and metallic tin powder (Sigma-Aldrich) in toluene according to a literature procedure [6]. The base hydrolysis of di(4-methylbenzyl)tin dichloride using 10% sodium hydroxide solution (Merck) afforded the di(4-methylbenzyl)tin oxide. Diisopropylthiocarbamoylthioacetic acid was synthesized from diisopropylamine (Merck), carbon disulfide (Merck) and chloroacetic acid (Sigma-Aldrich) according to a literature procedure [7]. Di(4-methylbenzyl)tin oxide (0.75 g, 2.0 mmol) and diisopropylthiocarbamoylthioacetic acid (0.94 g, 2.0 mmol) were heated in 95% ethanol (100 mL) for 1 h until the oxide dissolved. After filtration, the filtrate was evaporated slowly until colourless crystals were formed. Yield: 0.75 g (20%). M. pt: 424–426 K. Calcd. for $C_{100}H_{136}O_{10}N_4S_8Sn_4$: C 51.19; H 6.00; N 2.45%. Found: C 50.91; H 5.88; N 2.66%. IR (cm⁻¹) 503 (*m*) $\nu(\text{Sn}-\text{O})$, 639 (*m*) $\nu(\text{Sn}-\text{O}-\text{Sn})$, 1411, 1378 (*s*) $\nu_{\text{sym}}(\text{COO})$, 1662, 1597 (*s*) $\nu_{\text{asym}}(\text{COO})$.

Experimental details

The C-bound H atoms were geometrically placed (C–H = 0.93–0.98 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2-1.5U_{\text{eq}}(\text{C})$. Owing to poor agreement, five reflections, i.e. (0 0 1), (16 7 1), (1 1 1), (17 7 2) and (15 7 0), were omitted from the final cycles of refinement.

*Corresponding author: Edward R.T. Tiekink, Research Centre for Crystalline Materials, School of Science and Technology, Sunway University, 47500 Bandar Sunway, Selangor Darul Ehsan, Malaysia, e-mail: edwardt@sunway.edu.my, Edward.Tiekink@gmail.com. <https://orcid.org/0000-0003-1401-1520>

See Mun Lee and Kong Mun Lo: Research Centre for Crystalline Materials, School of Science and Technology, Sunway University, 47500 Bandar Sunway, Selangor Darul Ehsan, Malaysia

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} [*] / <i>U</i> _{eq}
Sn1	0.06743(2)	0.58266(2)	0.45775(2)	0.01558(7)
Sn2	0.15015(2)	0.36975(2)	0.34440(2)	0.01747(7)
S1	0.42834(8)	0.59921(9)	0.17472(7)	0.0328(2)
S2	0.26061(8)	0.78921(9)	0.07176(7)	0.0365(2)
S3	0.03115(8)	0.09307(7)	0.33558(6)	0.02718(19)
S4	−0.16901(8)	0.30322(9)	0.31429(7)	0.0360(2)
O1	0.05008(18)	0.44303(18)	0.44150(14)	0.0181(5)
O2	0.19911(18)	0.50238(19)	0.32557(15)	0.0203(5)
O3	0.3275(2)	0.4344(2)	0.22008(17)	0.0320(6)
O4	0.0680(2)	0.2578(2)	0.38506(18)	0.0297(6)
O5	−0.0586(2)	0.2745(2)	0.50262(16)	0.0236(5)
N1	0.4801(2)	0.7224(3)	0.02918(19)	0.0278(7)
N2	−0.0466(3)	0.1949(3)	0.1902(2)	0.0387(8)
C1	0.2792(3)	0.5079(3)	0.2607(2)	0.0214(7)
C2	0.2993(3)	0.6088(3)	0.2414(2)	0.0257(7)
H2A	0.297168	0.624248	0.294855	0.031*
H2B	0.240496	0.670124	0.213068	0.031*
C3	0.3925(3)	0.7092(3)	0.0837(2)	0.0244(7)
C4	0.4710(3)	0.8154(4)	−0.0479(3)	0.0363(9)
H4	0.546978	0.804128	−0.074974	0.044*
C5	0.4174(4)	0.8129(4)	−0.1156(3)	0.0470(12)
H5A	0.455285	0.742366	−0.130359	0.071*
H5B	0.421516	0.869466	−0.165831	0.071*
H5C	0.341579	0.825255	−0.093244	0.071*
C6	0.4218(4)	0.9262(4)	−0.0239(3)	0.0431(11)
H6A	0.345675	0.943736	0.000702	0.065*
H6B	0.427534	0.981909	−0.074382	0.065*
H6C	0.461102	0.922835	0.017059	0.065*
C7	0.5963(3)	0.6458(3)	0.0413(2)	0.0320(9)
H7	0.592182	0.590591	0.094576	0.038*
C8	0.6515(4)	0.5849(5)	−0.0301(3)	0.0551(14)
H8A	0.610328	0.545296	−0.031057	0.083*
H8B	0.724958	0.533998	−0.020643	0.083*
H8C	0.654391	0.636536	−0.083933	0.083*
C9	0.6587(4)	0.7049(5)	0.0541(4)	0.0548(13)
H9A	0.672227	0.753785	0.001042	0.082*
H9B	0.727629	0.651701	0.072009	0.082*
H9C	0.615653	0.746377	0.097228	0.082*
C10	−0.0019(3)	0.2298(3)	0.4408(2)	0.0199(7)
C11	−0.0161(3)	0.1306(3)	0.4375(2)	0.0267(7)
H11A	−0.093701	0.144209	0.453912	0.032*
H11B	0.022869	0.068377	0.479896	0.032*
C12	−0.0658(3)	0.2026(3)	0.2714(3)	0.0288(8)
C13	−0.1116(4)	0.2826(4)	0.1233(3)	0.0440(11)
H13	−0.079378	0.255229	0.070405	0.053*
C14	−0.0954(5)	0.3886(4)	0.1040(3)	0.0526(13)
H14A	−0.137608	0.427029	0.149119	0.079*
H14B	−0.119514	0.434147	0.050496	0.079*
H14C	−0.018788	0.371765	0.100293	0.079*
C15	−0.2307(4)	0.2978(5)	0.1402(4)	0.0631(15)
H15A	−0.235145	0.227269	0.161622	0.095*
H15B	−0.261102	0.333994	0.087830	0.095*
H15C	−0.271560	0.341978	0.181954	0.095*
C16	0.0427(5)	0.1001(4)	0.1572(3)	0.0601(16)
H16	0.077435	0.047404	0.206563	0.072*

Table 2 (continued)

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} [*] / <i>U</i> _{eq}
C17	0.1312(5)	0.1337(6)	0.0943(4)	0.080(2)
H17A	0.101694	0.183376	0.043673	0.119*
H17B	0.192248	0.069402	0.079162	0.119*
H17C	0.155740	0.169417	0.120375	0.119*
C18	0.0000(6)	0.0414(5)	0.1230(4)	0.083(2)
H18A	−0.057465	0.024947	0.164866	0.124*
H18B	0.058865	−0.025889	0.111039	0.124*
H18C	−0.028785	0.087533	0.071155	0.124*
C19	−0.0052(3)	0.6997(3)	0.3498(2)	0.0219(7)
H19A	0.050772	0.723579	0.311100	0.026*
H19B	−0.022505	0.661147	0.320068	0.026*
C20	−0.1074(3)	0.8019(3)	0.3631(2)	0.0204(7)
C21	−0.0998(3)	0.8989(3)	0.3590(2)	0.0240(7)
H21	−0.030880	0.899272	0.353335	0.029*
C22	−0.1938(3)	0.9961(3)	0.3634(2)	0.0285(8)
H22	−0.186507	1.060318	0.360261	0.034*
C23	−0.2981(3)	0.9982(3)	0.3724(2)	0.0297(8)
C24	−0.3055(3)	0.9013(3)	0.3778(2)	0.0319(8)
H24	−0.374710	0.900862	0.385022	0.038*
C25	−0.2125(3)	0.8038(3)	0.3729(2)	0.0252(7)
H25	−0.220287	0.739835	0.376170	0.030*
C26	−0.3998(4)	1.1045(4)	0.3771(3)	0.0475(12)
H26A	−0.462087	1.096027	0.369836	0.071*
H26B	−0.388025	1.163411	0.332436	0.071*
H26C	−0.413611	1.121115	0.431909	0.071*
C27	0.2161(3)	0.4893(3)	0.5129(2)	0.0215(7)
H27A	0.200636	0.501921	0.570031	0.026*
H27B	0.234015	0.411858	0.519487	0.026*
C28	0.3165(3)	0.5108(3)	0.4661(2)	0.0216(7)
C29	0.4042(3)	0.4359(3)	0.4217(2)	0.0244(7)
H29	0.400849	0.370730	0.422148	0.029*
C30	0.4958(3)	0.4572(3)	0.3772(2)	0.0261(7)
H30	0.552788	0.406344	0.347780	0.031*
C31	0.5042(3)	0.5531(3)	0.3756(2)	0.0245(7)
C32	0.4193(3)	0.6256(3)	0.4220(2)	0.0262(7)
H32	0.424730	0.688937	0.423619	0.031*
C33	0.3261(3)	0.6058(3)	0.4663(2)	0.0236(7)
H33	0.269733	0.656358	0.496266	0.028*
C34	0.6023(3)	0.5793(4)	0.3252(3)	0.0338(9)
H34A	0.667105	0.512118	0.324875	0.051*
H34B	0.612363	0.625563	0.351491	0.051*
H34C	0.589740	0.616719	0.267361	0.051*
C35	0.0938(3)	0.4180(3)	0.2241(2)	0.0253(7)
H35A	0.157492	0.399247	0.179796	0.030*
H35B	0.052988	0.375255	0.227274	0.030*
C36	0.0219(3)	0.5382(3)	0.1976(2)	0.0249(7)
C37	0.0662(3)	0.6143(3)	0.1495(2)	0.0276(8)
H37	0.141704	0.590471	0.130369	0.033*
C38	−0.0006(3)	0.7255(3)	0.1297(2)	0.0318(8)
H38	0.031229	0.775083	0.097883	0.038*
C39	−0.1138(3)	0.7649(3)	0.1560(3)	0.0341(9)
C40	−0.1582(3)	0.6886(4)	0.2019(3)	0.0344(9)
H40	−0.234062	0.712510	0.219008	0.041*
C41	−0.0917(3)	0.5761(3)	0.2231(2)	0.0292(8)
H41	−0.123594	0.526435	0.254481	0.035*
C42	−0.1848(4)	0.8859(4)	0.1388(3)	0.0505(12)

Table 2 (continued)

Atom	<i>x</i>	<i>y</i>	<i>z</i>	U_{iso}^*/U_{eq}
H42A	−0.255132	0.897749	0.128725	0.076*
H42B	−0.149809	0.923386	0.089199	0.076*
H42C	−0.194958	0.914038	0.187564	0.076*
C43	0.2986(3)	0.2319(3)	0.3784(3)	0.0279(8)
H43A	0.360903	0.252456	0.354138	0.033*
H43B	0.294110	0.210778	0.440337	0.033*
C44	0.3154(3)	0.1361(3)	0.3456(2)	0.0257(7)
C45	0.3446(3)	0.1391(3)	0.2577(3)	0.0326(9)
H45	0.356714	0.199449	0.219999	0.039*
C46	0.3557(4)	0.0538(3)	0.2257(3)	0.0374(10)
H46	0.375480	0.057846	0.166963	0.045*
C47	0.3380(3)	−0.0371(3)	0.2791(3)	0.0352(9)
C48	0.3114(4)	−0.0413(3)	0.3668(3)	0.0363(9)
H48	0.300875	−0.102408	0.404493	0.044*
C49	0.3004(3)	0.0441(3)	0.3993(3)	0.0325(9)
H49	0.282499	0.039028	0.458175	0.039*
C50	0.3471(5)	−0.1277(4)	0.2428(3)	0.0539(13)
H50A	0.322138	−0.096562	0.188139	0.081*
H50B	0.302517	−0.165096	0.281434	0.081*
H50C	0.422555	−0.179160	0.235740	0.081*

Comment

Organotin(IV) carboxylates have been widely investigated because of their structural diversity [8] and their potential biological properties, in particular anti-tumour potential [9, 10]. In addition, organotin compounds containing dithiocarbamate anions have also received much attention due to their promising anti-fungal, anti-bacterial and anti-tumour activities [11, 12]. In continuation of our efforts in exploring the coordination chemistry and anti-proliferative activities of organotin carboxylates, herein the synthesis and structural features of an organotin compound with a carboxylate ligand containing a dithiocarbamate fragment are described.

The title compound is a centrosymmetric, tetra-nuclear species shown in the figure (50% displacement ellipsoids; unlabelled atoms are related by the symmetry operation (i): $-x, 1-y, 1-z$). The molecule is constructed about a central Sn_2O_2 core with $Sn1-O1$, $O1^i$ bond lengths of 2.173(2) and 2.050(2) Å, respectively. Connected to this ring is the exocyclic Sn_2 atom [$Sn2-O1=2.026(2)$ Å]. The additional link between the endocyclic- $Sn1$ and exocyclic- Sn_2 atoms is provided by an almost symmetrically bridging carboxylate ligand [$Sn2-O4=2.164(2)$ Å and $Sn1-O5=2.237(2)$ Å]. The exocyclic- Sn_2 atom is also coordinated by a monodentate carboxylate ligand [$Sn2-O2=2.151(2)$ Å] with the $Sn_2 \cdots O3$ separation of 3.012(3) Å considered too long for a significant bonding interaction. There is some evidence for the $O2$ atom semi-bridging the $Sn1$ atom as the $Sn1 \cdots O2$ separation is 2.628(2) Å. The five-coordinate geometries for each of the $Sn1$ and Sn_2 atoms is completed by two carbon atoms derived from

the benzyl substituents. To a first approximation, the C_2O_3 donor sets define distorted trigonal bipyramidal geometries with $O1-Sn1-O5$ [168.41(9)°] and $O2-Sn2-O4$ [165.91(9)°] axial angles. Each of the carboxylate ligands is twisted as seen in the dihedral angles between the CO_2 and CS_2 residues of 70.0(4) and 81.3(3)° for the $O2$ - and $O4$ -carboxylate ligands, respectively.

Molecules stack in columns along the *a* axis being sustained by tolyl- $Me-C-H \cdots S$ (thione) [$C34-H34a \cdots S4^{ii}$: $H34a \cdots S4^{ii}=2.79$ Å, $C34 \cdots S4^{ii}=3.741(5)$ Å with an angle at $H34a=169^\circ$ for symmetry operation (ii) $1+x, y, z$] and $\pi \cdots \pi$ interactions between centrosymmetrically related tolyl moieties [inter-centroid $Cg(C28-C33) \cdots Cg(C28-C33)^{iii}$ separation = 3.709(2) Å for symmetry operation (iii): $1-x, 1-y, 1-z$]. The chains assemble in the three-dimensional architecture without directional interactions between them.

Tetranuclear clusters of the general formula $\{[R_2SnX]_2O\}_2$ are common hydrolysis products of diorganotin species [13]. In keeping with this observation, there are at least five structures of this type containing dithiocarbamate-functionalized carboxylates, *i.e.* $R'_2NC(=S)SCH_2CO_2^-$. The three structures with $R=n$ -Bu and $R'=Et$ [14], $R'_2=(CH_2)_4$ [15] and $R'_2=(CH_2)_5$ [16] adopt the same structural motif as the title structure. The two structures with $R=n$ -Bu and $R'=Me$ [17] and $R=n$ -Oct and $R'_2=(CH_2CH_2)_2O$ [18] adopt essentially the same motif but the carboxylate bridge involves one oxygen atom only. Both motifs have ample precedents in the crystallographic literature of the diorganotin bis(carboxylates) [8].

Acknowledgements: Sunway University is thanked for supporting studies in organotin chemistry.

References

1. Bruker. APEX2 and SAINT. Bruker AXS Inc., Madison, WI, U.S.A. (2008).
2. Sheldrick, G. M.: SADABS. University of Göttingen, Germany (1996).
3. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr. A* **64** (2008) 112–122.
4. Sheldrick, G. M.: Crystal structure refinement with SHELXL. *Acta Crystallogr. C* **71** (2015) 3–8.
5. Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Crystallogr.* **45** (2012) 849–854.
6. Sisido, K.; Takeda, Y.; Kinugawa, Z.: Direct synthesis of organotin compounds I. di- and tribenzyltin chlorides. *J. Am. Chem. Soc.* **83** (1961) 538–541.
7. Nachmias, G.: Sur quelques nouveaux derives de lacide dithiocarbamique *N*-disubstitues. *C. r. hebdomadaire des seances Acad. Sci.* **232** (1951) 1118–1119.
8. Tieckink, E. R. T.: Structural chemistry of organotin carboxylates: a review of the crystallographic literature. *Appl. Organomet. Chem.* **5** (1991) 1–23.

9. Gielen, M.: Organotin compounds and their therapeutic potential: a report from the Organometallic Chemistry Department of the Free University of Brussels. *Appl. Organomet. Chem.* **16** (2002) 481–494.
10. Anasamy, T.; Thy, C. K.; Lo, K. M.; Chee, C. F.; Yeap, S. K.; Kamalidehghan, B.; Chung, L. Y.: Tribenzyltin carboxylates as anticancer drug candidates: effect on the cytotoxicity, motility and invasiveness of breast cancer cell lines. *Eur. J. Med. Chem.* **125** (2017) 770–783.
11. Tiekink, E. R. T.: Tin dithiocarbamates: applications and structures. *Appl. Organomet. Chem.* **22** (2008) 533–550.
12. Khan, N.; Farina, Y.; Lo, K. M.; Rajab, N. F.; Awang, N.: Synthesis, spectral characterization, X-ray studies and *in vitro* cytotoxic activities of triorganotin(IV) derivatives of *p*-substituted *N*-methylbenzylaminedithiocarbamates. *J. Mol. Struct.* **1076** (2014) 403–410.
13. Dakternieks, D.; Jurkschat, K.; van Dreumel, S.; Tiekink, E. R. T.: Molecular dynamics within diorganotin systems: solution and solid state studies of new mixed distannoxane dimers [tBu₂(Cl)SnOSn(Cl)R₂]₂. *Inorg. Chem.* **36** (1997) 2023–2029.
14. Ng, S. W.; Kumar Das, V. G.: Tetrabutylbis[(*N,N*-diethylthiocarbamoylthio)acetato] distannoxane dimer. *Acta Crystallogr.* **C51** (1995) 1774–1776.
15. Yin, H.-D.; Xue, S.-C.; Wang, Q.-B.: Synthesis, characterization and crystal structure of the dimeric organotin compound: {[*n*-Bu₂Sn(O₂CCH₂CS₂NC₄H₈)]₂O}₂. *Chin. J. Inorg. Chem.* **20** (2004) 421–425.
16. Yin, H.-D.; Xue, S.-C.; Wang, Q.-B.: Synthesis, characterization of di-*n*-butyltin compounds with (thiocarbamoylthio)acetic acid and crystal structure of {[*n*-Bu₂Sn(O₂CCH₂CS₂NC₅H₁₀)]₂O}₂·2 H₂O and *n*-Bu₂Sn(O₂CCH₂CS₂NC₅H₁₀)₂. *Indian J. Chem., Sect. B* **44** (2005) 1040–1045.
17. Yin, H.-D.; Xue, S.-C.; Liu, G.-F.: Synthesis and crystal structure of bis{oxo-bis[*N,N*-dimethylthiocarbamoylthioacetato-di-*n*-butyltin(IV)]} and bis{oxo-bis[*N,N*-diethylthiocarbamoylthioacetato di-*n*-butyltin(IV)]}. *Acta Chim. Sinica* **62** (2004) 603–609.
18. Yin, H.-D.; Gao, Z.-J.; Li, G.; Xu, H.-L.; Hong, M.: Synthesis and crystal structure of the dimeric organotin complex {[(*n*-C₈H₁₇)₂Sn(O₂CCH₂CS₂NC₄H₈O)]₂O}₂. *Chin. J. Inorg. Chem.* **22** (2006) 157–160.